

ON THE CONSTELLATIONS OF WEIERSTRASS POINTS

FERNANDO TORRES

ABSTRACT. We prove that the constellation of Weierstrass points characterizes the isomorphism-class of double coverings of curves of genus large enough.

1. Let X be a projective, irreducible, non-singular algebraic curve defined over an algebraically closed field k of characteristic p . Let $n \geq 1$ be an integer and C_X a canonical divisor of X . The pluricanonical linear system $|nC_X|$ defines a nondegenerate morphism

$$\pi_n : X \rightarrow \mathbb{P}^{N(n)},$$

where $N(1) = g - 1$, and $N(n) = (2n - 1)(g - 1) - 1$ for $n \geq 2$. To any $P \in X$ we then associate the sequence of multiplicities

$$\{v_P(\pi_n^*(H)) : H \text{ hyperplane} \subseteq \mathbb{P}^{N(n)}\} = \{\epsilon_0(P) < \epsilon_1(P) < \dots < \epsilon_{N(n)}(P)\}.$$

Such a sequence is the same for all but finitely many points (cf. [F-K, III.5], [Lak, Prop.3], [S-V, §1]). These finitely many points are the so called n -Weierstrass points of X . There exists a divisor $\mathcal{W}_n$ on X whose support is the set of n -Weierstrass points and satisfies the property below. Let denote by $v_P(\mathcal{W}_n)$ the coefficient of $\mathcal{W}_n$ in P (called the n -Weierstrass weight at P). Then

$$(1) \quad \omega_n := \deg(\mathcal{W}_n) = \sum_P v_P(\mathcal{W}_n) = \sum_{i=0}^{N(n)} \epsilon_i(2g - 2) + n(2g - 2)(N(n) + 1),$$

where $\epsilon_0 < \epsilon_1 < \dots < \epsilon_{N(n)}$ denotes the sequence at a generic point ([F-K, III.5], [Lak, Thm.6], [S-V, p.6]). One has $\epsilon_i(P) \geq \epsilon_i$ for each i and for each P ([F-K, III.5], [Lak, Prop.3], [S-V, p.5]).

Let $P_1, \dots, P_{\omega_n}$ be the n -Weierstrass points (counted with multiplicity according to their n -Weierstrass weights).

Definition. The orbit $\mathcal{O}_n(X)$ of $(\pi_n(P_1), \dots, \pi_n(P_{\omega_n})) \in (\mathbb{P}^{N(n)})^{\omega_n}$ under the action of the product of the symmetric group S_{ω_n} and the projective linear group $PGL(N(n) + 1)$ is called the constellation of n -Weierstrass points of X . $\mathcal{O}_1(X)$ is called the constellation of Weierstrass points of X .

Let X, X_1 be curves as above of genus g . Then clearly $\mathcal{O}_n(X) = \mathcal{O}_n(X_1)$ if X is isomorphic to X_1 . In this note we are interested in the converse:

$$(P) \quad \mathcal{O}_n(X) = \mathcal{O}_n(X_1) \text{ for some } n \geq 1 \Rightarrow X \text{ is isomorphic to } X_1 ?$$

This problem was studied by Pflaum [Pf] for $p = 0$. He showed that (P) is true in the following cases :

- (i) If $n \geq 2$;
- (ii) If $n = 1$, and $2 \leq g \leq 15$;

The author is supported by a grant from the International Atomic Energy Agency and UNESCO.

- (iii) If $n = 1$, and X and X_1 are γ -hyperelliptic curves (that is double coverings of curves of genus γ) with $g \geq 2$ and $\gamma \in \{0, 1, 2\}$.

For the cases $n = 2, g = 2$; $n = 1, g = 3, 4$; and $n = 1, X, X_1$ hyperelliptic curves the proof is given by a direct application of the definition above. For the remaining cases Pflaum used the following argument. By means of a lower bound on the number W_n of n -Weierstrass points, he stated (Corollary 2.6, loc. cit.) a sufficient condition to have an affirmative answer to the problem (this condition holds regardless the characteristic of k). Define the number $N(g, n)$ by $N(6, 1) = 25$, $N(g, 1) = \max\{3g + 6, 4g - 4\}$ for $g \neq 6$, and $N(g, n) = 4n(g - 1)$ for $n \geq 2$. Pflaum then showed that (P) is true provided

$$(2) \quad W_n > N(g, n).$$

A way of getting a lower bound on W_n is by bounding from above $v_P(\mathcal{W}_n)$, and then by using (1). One has

$$(3) \quad v_P(\mathcal{W}_n) \geq \sum_{i=1}^{N(n)} (\epsilon_i(P) - \epsilon_i)$$

for each P , and equality holds if

$$\det \left(\begin{pmatrix} \epsilon_i(P) \\ \epsilon_j \end{pmatrix} \right) \not\equiv 0 \pmod{p}$$

([Lak, p.239], [S-V, Thm.1.5]). A curve X is called *classical* (with respect to the pluricanonical linear system $|nC_X|$) if $\epsilon_i = i$ for each i . This is the case if $p = 0$ or $p > n(2g - 2)$, and here one also has the equality in (3) for each P ([F-K, III.5], [S-V, Corollary 1.8], [Lak, Thm.11]).

Let X and X_1 be classical curves of genus g , and suppose that we have equality in (3) for each point of X and X_1 . If $n = 1$ and $5 \leq g \leq 15$, Pflaum showed (2) by direct computations. If $n \geq 2$, then Homma and Ommori [H-O] noticed that $v_P(\mathcal{W}_n) \leq g(g + 1)/2$. Hence by means of (1) Pflaum obtained (2). Now if X and X_1 are γ -hyperelliptic curves of genus large enough with $\gamma \in \{1, 2\}$ and if $p = 0$ or $p > 2g - 2$, by using results due to Garcia [G] and Kato [K] Pflaum can bound from above $v_P(\mathcal{W}_1(P))$ and obtain (2).

The aim of this note is to extend Pflaum's result (iii) above to γ -hyperelliptic curves of genus large enough ($\gamma \geq 3$) and whenever $p = 0$ or $p > 2g - 2$. We will show that such curves satisfy (2) and to do that we use some results concerning Weierstrass weights in [To1] and [To2]. We show

Theorem. *Let X and X_1 be γ -hyperelliptic curves of genus $g \geq 9\gamma - 17 + \frac{43\gamma - 20}{2\gamma^2 + \gamma - 1}$ with $\gamma \geq 3$. Assume that $p = 0$ or $p > 2g - 2$. Then X and X_1 are isomorphic provided $\mathcal{O}_1(X) = \mathcal{O}_1(X_1)$.*

In general one cannot expect to fulfil condition (2) for $0 < p \leq 2g - 2$, because in this case there exist curves with small number of Weierstrass points. For instance there exist curves with just one Weierstrass point (see [Lak, §6]).

2. Let

$$\pi : X \rightarrow \tilde{X}$$

be a double covering of curves of genus g and $\gamma \geq 3$ respectively. Let $P \in X$, and set $w(P) := v_P(\mathcal{W}_1)$. The key point of the proof is the fact that we can bound from above $w(P)$ by considering the following three cases:

- (I) P is a ramified point of π such that $\pi(P)$ is a Weierstrass point of $\tilde{X}$.
- (II) P is a ramified point of π such that $\pi(P)$ is not a Weierstrass point of $\tilde{X}$.
- (III) P is not a ramified point of π .

Lemma. *Let X be a γ -hyperelliptic curve of genus g ($\gamma \geq 3$). Let assume that $p = 0$ or $p > 2g - 2$, and let $P \in X$.*

- (i) *If P is as in (I), then*

$$w(P) \leq c_1 := \binom{g-2\gamma}{2} + 2\gamma^2.$$

- (ii) *If P is as in (II), then*

$$w(P) \leq c_2 := \binom{g-2\gamma}{2} + 4\gamma - 4.$$

- (iii) *If P is as in (III) and $g \geq 2\gamma$, then*

$$w(P) \leq c_3 := \max\{(2\gamma - 1)g - (\gamma - 1)(2\gamma + 1), 2\gamma g - 2\gamma(4\gamma - 1)\}.$$

3. Proof of the Theorem. Since $g \geq 2\gamma + 2$, every ramified point of π is a Weierstrass point (see §4). Let t denote the number of points of type (I). Then by the lemma we have:

$$\begin{aligned} \deg(\mathcal{W}_1) &= g^3 - g \leq tc_1 + (2g - 4\gamma + 2 - t)c_2 + (W_1 - 2g + 4\gamma - 2)c_3 \\ &= (c_1 - c_2)t + (2g - 4\gamma + 2)c_2 + (W_1 - 2g + 4\gamma - 2)c_3. \end{aligned}$$

Then by noticing that $t \leq \min\{\gamma^3 - \gamma, 2g - 4\gamma + 2\}$, we find

$$(4) \quad (W_1 - 2g + 4\gamma - 2)c_3 \geq 6\gamma g^2 - 16\gamma^2 g + 16\gamma^3 - 4\gamma^2 - 2\gamma.$$

If $g \geq 6\gamma^2 - \gamma + 1$, then $c_3 = 2\gamma(g - 4\gamma + 1)$ and from (4) we get

$$W_1 \geq 5g - 1 + \frac{24\gamma^2 - 10\gamma - 4}{g - 4\gamma + 1} > N(g, 1).$$

Now suppose that $c_3 = (2\gamma - 1)g - (2\gamma^2 - \gamma - 1)$. From (4) we find that $W_1 > N(g, 1)$ if

$$(2\gamma - 1)(2\gamma + 2)g - (36\gamma^3 - 50\gamma^2 + 34\gamma - 6) + \frac{24\gamma^5 - 40\gamma^4 + 22\gamma^3 + 4\gamma}{(2\gamma - 1)g - (2\gamma^2 - \gamma - 1)} > 0.$$

This is satisfied provided $g \geq 9\gamma - 17 + \frac{43\gamma - 20}{2\gamma^2 + \gamma - 1}$.

4. Proof of the Lemma. First we recall some properties of Weierstrass semigroups. Let $P \in X$. In the case of 1-Weierstrass points, the set

$$G(P) := \{\epsilon_i(P) + 1 : 0 \leq i \leq g - 1\}$$

is the complement (or the gaps) of a semigroup $H(P)$, the so called Weierstrass semigroup at P . $H(P)$ looks like

$$H(P) = \{0 < m_1(P) < \dots < m_g(P) = 2g < 2g + 1 < \dots\},$$

and it is satisfied the following property ([B], [Oliv, Thm. 1.1(ii)]). Let $\ell_i(P) := \epsilon_i(P) + 1$. Then

$$(5) \quad \ell_i(P) \leq 2i - 2, \quad \text{for} \quad i = 2, \dots, g - 1, \quad \ell_g(P) \leq 2g - 1,$$

provided $m_1(P) \geq 3$. Then if X is classical and if we have equality in (3), $w(P)$ can be computed by the formula

$$(6) \quad w(P) = \frac{3g^2 + g}{2} - \sum_{m \in H(P), m \leq 2g} m.$$

Now in case of γ -hyperelliptic curves, P a ramified point of π and $p > 2$, $H(P)$ fulfil the following properties ([To1, Lemma 3.4]):

- (A) $\gamma = \#\{\ell \in G(P) : \ell \text{ even}\}$.
- (B) $H(\pi(P)) = \{\frac{h}{2} : h \in H(P), h \text{ even}\}$.

(Note that property (B) implies $h \leq 2\gamma + 2$ for $h \in H(P)$, h even. In particular if X is classical and $g \geq 2\gamma + 2$, then each ramified point of π is a Weierstrass point of X .)

Proof of (i). Follows from property (A) above and [To2, Lemma 3.1.2(ii)].

Proof of (ii). Let $P \in X$ be as in (II). Since $p = 0$ or $p > 2g - 2 > 2\gamma - 2$, then $\tilde{X}$ is also a classical curve. Thus from properties (A) and (B) we have that all the even positive non-gaps of H belong to the following set:

$$\{2\gamma + 2i : i \in \{1, \dots, g - \gamma\}\}.$$

Hence

$$(*) \quad \sum_{h \in H(P), h \text{ even}, h \leq 2g} h = g^2 + g - \gamma^2 - \gamma.$$

Let denote by $u_\gamma < \dots < u_1$ the γ odd non-gaps at P in $[1, 2g - 1]$. According to (6) and (*), an upper bound for $w(P)$ corresponds to a lower bound for $\sum_{i=1}^\gamma u_i$. By [To1, Lemma 2.1], $u_\gamma \geq 2g - 4\gamma + 1$. If $u_\gamma \geq 2g - 2\gamma - 1$, any odd number in $[2g - 2\gamma + 1, 2g - 1]$ could be an odd non-gap at P . Hence in this case we have

$$\sum_{i=1}^\gamma u_i \geq 2\gamma g - \gamma^2 - 2\gamma.$$

If $u_\gamma \leq 2g - 2\gamma - 3$ (then $\gamma \geq 2$), it is easy to see that the minimum for $\sum_{i=1}^\gamma u_i$ is reached for the sequence $2g - (2i + 5), 2g - 1, i = 1, \dots, \gamma - 1$. Hence in this case we have

$$\sum_{i=1}^\gamma u_i \geq 2\gamma g - \gamma^2 - 4\gamma + 4.$$

Then since $\gamma > 1$ from (*), the last inequality and (6) we obtain (ii).

Proof of (iii). Let $P \in X$ and suppose that P is not a ramified point of π .

Claim. Let h be a non-gap at P . Then $h \geq g - 2\gamma + 1$.

Remark. Let $f \in k(X)$ and denote by $O(f)$ the degree of f . Then $O(f)$ is even provided $O(f) < g + 1 - 2\gamma$ and $g \geq 4\gamma + 2$. For $p = 0$ this is a result due to Farkas [F, Thm.2(iii)] (see also [F-K, Thm.V.1.9], Accola [A, Lemma 4]) and in general is due to Stichtenoth [Sti, Satz 2]. The claim follows from this result but with an extra hypothesis on g . We will see that in the case that f has just one pole one can avoid such a hypothesis. The claim is a particular case of [To1, Corollary 3.3(ii)], and for the sake of completeness we state a proof of it.

Proof. (Claim.) Suppose that $h < g - 2\gamma + 1$. Consider $K' := k(\tilde{X}).k(f)$, with $\text{div}_\infty(f) = hP$. Then by Castelnuovo's inequality concerning subfields of $k(X)$ (see [C], [Sti1]) we must have $K' = k(\tilde{X})$. Thus there exists $\tilde{f} \in K(\tilde{X})$ such that $f = \tilde{f} \circ \pi$ and we would have that P is a totally ramified point of π , a contradiction. $\square$

Let $g \geq 2\gamma$. We have

$$(7) \quad w(P) = \sum_{i=g-2\gamma+1}^g (\ell_i - i),$$

and then we consider two cases:

- (a) There exists $\ell \in G(P) \cap [g - 2\gamma + 1, g]$.
- (b) $[g - 2\gamma + 1, g] \cap \mathbb{N} \subseteq H(P)$.

In the first case we have $\ell_{g-2\gamma+1} = g - 2\gamma + j$ with $j \in \{1, \dots, \gamma\}$. Then from (7) and (5) we get

$$w(P) \leq (2\gamma - 1) + \sum_{i=g-2\gamma+2}^{g-1} (i - 2) + (g - 1) = (2\gamma - 1)g - (\gamma - 1)(2\gamma + 1).$$

In the second case, due to the semigroup property of $H(P)$, $w(P)$ reaches its maximum whenever $G(P) = \{1, \dots, g - 2\gamma, 2g - 6\gamma + 2, \dots, 2g - 4\gamma + 1\}$. Then from (7) we find

$$w(P) \leq 2\gamma g - 2\gamma(4\gamma - 1).$$

This finish the proof of the lemma.

REFERENCES

- [A] Accola, R.D.M.: *Strongly branched coverings of closed Riemann surfaces*, Proc. Amer. Math. Soc. **26** (1970), 315–322.
- [B] Buchweitz, R.O.: “Über deformationem monomialer kurvensingularitäten und Weierstrasspunkte auf Riemannschen flächen”, Thesis, Hannover 1976.
- [C] Castelnuovo, G.: *Sulle serie algebriche di gruppi di punti appartenenti ad una curva algebrica*, Rendiconti della Reale Accademia dei Lincei (5) **15** (1906), 337–344. Reprinted in Memoria scelta, Zanichelli, Bologna, 509–517, 1937.
- [F] Farkas, H.M.: *Remarks on automorphisms of compact Riemann surfaces*, Ann. of Math. Stud. **78** (1974), 121–144.
- [F-K] Farkas, H.M.; Kra, I.: “Riemann surfaces”, Grad. Texts in Math. **71** (second edition) Springer-Verlag 1992.
- [G] Garcia, A.: *Weights of Weierstrass points in double covering of curves of genus one or two*, Manuscripta Math. **55** (1986), 419–432.
- [H-O] Homma, M.; Ommori, S.: *On the weight of higher order Weierstrass points*, Tsukuba J. Math. **8** (1984), 189–198.
- [K] Kato, T.: *Non-hyperelliptic Weierstrass points of maximal weight*, Math. Ann. **239**, (1979), 141–147.
- [Lak] Laksov, D.: *Weierstrass points on curves*, Astérisque **87-88**, (Société Mathématique de France, Paris, 1981), 221–247.
- [Oliv] Oliveira, G.: *Weierstrass semigroups and the canonical ideal of non-trigonal curves*, Manuscripta Math. **71** (1991), 431–450.
- [Pf] Pflaum, U.: *The canonical constellations of k -Weierstrass points*, Manuscripta Math. **59** (1987), 21–34.
- [Sti] Stichtenoth, H.: *s -Erweiterungen algebraischer Funktionenkörper*, Arch. Math. **43** (1984), 27–31.
- [Sti1] Stichtenoth, H.: *Die Ungleichung von Castelnuovo*, J. Reine Angew. Math. **348** (1984), 197–202.

- [S-V] Stöhr, K.O.; Voloch, J.F.: *Weierstrass points and curves over finite fields*, Proc. London Math. Soc. (3), **52** (1986), 1–19.
- [To1] Torres, F.: *On certain N -sheeted coverings of curves and numerical semigroups which cannot be realized as Weierstrass semigroups*, Comm. Algebra **23** (11) (1995), 4211–4228.
- [To2] Torres, F.: *Remarks on numerical semigroups*, alg-geo e-print 9512012.

ICTP, MATHEMATICS SECTION, P.O. BOX 586, 34100, TRIESTE - ITALY
E-mail address: `feto@ictp.trieste.it`